

FREDUN PHARMACEUTICALS LTD

VETERINARY PRODUCT LIST

ANIMALS / POULTRY MEDICINES & SUPPLEMENTS					
GROWTH PROMOTOR					
Brand Name		Composition	Indications	Dosage	Presentation
GROWAY	Bacitracin Methylene Disalicylate 10% Animal Feed Premix	Each kg of GROWAY contains: Bacitracin Methylene Disalicylate 100 gms	Growth Promoter Improves FCR Gives better carcass quality Remarkable increase in egg production	For Growth Promotion 150-200 gms/ton of feed for broilers layers & breeders To optimize egg production 110-275 gms/ ton of feed	1 kg, 5 kgs , 10 kgs & 25 kg Bag
COLI-APPETI	Colistin Sulphate 10% Feed Premix	Each gm of Coli-appeti contains : Colistin Sulphate USP (equivalent to colistin base) 100 mg	Promotes Growth , Increases feed utilization, Antibacterial activity against Gram negative bacteria	For prophylaxis and growth promotion 50 gms/MT of feed For Treatment : 200 gms/ton of feed or Dissolve 2 gms in 15 to 20 Liters of drinking water for 75-100 birds for 3-5 day or, As Directed by Veterianian	50 gms & 200 packs 25 kg HDPE Laminated Bag
FATTY FACTS	Fats for Monogastric Animals (Swine, Poultry)	Total Fat Content : 99.0% Min Free Fatty Acid : 1-10% Iodine Value : 14-18	For economical high energy source & for Better growth	Add 2 – 5% of FATTY FACTS in the animal feed or consult a local veterinary nutritionist.	1 kg , 5 kgs, 10 kgs & 25 kgs Bag
ANTIBIOTICS/ ANTIBACTERIALS					
DOXYFRED-N	Doxycycline Hyclate + Neomycin Sulphate	Each g of DOXYFRED-N contains: Doxycycline Hyclate BP 115 mg eq. to Doxycycline 100 mg Neomycin Sulphate BP 143 mg eq. to Neomycin 100 mg	Fowl Cholera Mortality Complex CRD with E. Coli Non-specific diarrhoea Infection due to gram positive and gram negative bacteria	During stress : 1 gm in 10 ltrs of drinking water for 4 to 5 days for 60 growers or 45 adult birds. For Prevention of Early Chick Mortality: 1 gm in 10 ltrs of drinking water for 3 to 4 days for 100 chicks. In Severe acute Infections: 1 gm in 2 to 4 ltrs of drinking water for 4 to 5 days. For small/large animals (sheep, swine, dogs, horse, cattle etc.) 10 to 20 mg per kg body weight for 3 to 4 days or, till infection subsides.	1 Kg Aluminium Foil 25 kg HDPE Laminated Bag
DOXYFRED POWDER w/s	Doxycyclin Powder 20 %	Each 1 g contains Doxycycline Hyclate BP 221 mg eq to Doxycycline 200 mg	Gastrointestinal and respiratory infections caused by doxycycline sensitive micro-organisms like Bordetella, Campylobacter, Chlamydia, E. coli, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in calves, goats, poultry, sheep, and	For oral administration: Calves, goats and sheep Twice daily 5 g per 200 kg body weight for 3 - 5 days.	Sachet of 100 g and jar of 500 and 1000 g.
DOXYFRED - T	Doxycycline & Tylosin Tartrate W/S Powder For Animals & Poultry Use	Each gm contains Doxycycline Hyclate BP Equivalent to Doxycycline 100 mg Tylosin Tartrate BP 100 mg Excipients q.s.	It is effective against infections caused by micro-organisms in calves, poultry and pigs. Especially indicated for gastrointestinal and respiratory infections	For oral administration via feed or drinking water. Calves: 500 g per 400 calves of 50 kg body weight daily, for 3-5 days. Pigs: 100 g per 200-400 litres of drinking water daily, for 3-5 days Poultry: 100 g per 200-400 litres of drinking water daily, for 3-5 days For Prevention: Mix 500 gm per 1000 kg of feed. For Treatment: 1 kg per 1000 kg of feed, for 2-5 days. The above mentioned dosage for pigs and poultry can be doubled without any danger. Mix with feed, and use immediately. Medicated drinking water should be used within 24 hours.	100 gm , 227 gm & 1 kg pouch Pack.

ENROFRED	Enrofloxacin Oral Liquid 10%	Each ml of ENROFRED contains: Enrofloxacin 100 mg	For prevention of Chronic Respiratory Disease Broncho Pneumonia Cholera, typhoid, Paratyphoid, Colibacillosis, Colisepticaemia..etc	Withdrawal period: 15 days for Meat, 4 days for eggs.	1 Ltr HDPE Bottle 5 Ltrs Jar 25 Ltrs Drum
ERYFRED POWDER w/s	Erythromycin Thiocynate Powder 20 %	Each 100 g contains: Erythromycin Thiocynate 20% (Erythromycin 20 g Base)	ERYFRED POWDER W/S is indicated for the treatment of Chronic Respiratory Disease (CRD) caused by Mycoplasma infections, and for the treatment of infectious (respiratory) diseases due to erythromycin- sensitive bacteria such as Staphylococcus aureus (e.g. infectious arthritis), Haemophilus paragallinarum (e.g. infectious coryza) and Pasteurella multocida (e.g. pasteurellosis; pneumonia) in poultry.	For oral administration, Poultry: 100 g per 100 liter drinking water for 3-5 days. For meat: 3 days, For eggs: 6 days,	1 kg pouch , 25 kg HDPE laminated bag
FRAMOX- C	Amoxycillin & Colistin Sulphate Powder For Animals & Poultry Use	Each gm. contains Amoxycillin Trihydrate BP Equivalent to Amoxycillin 200 mg Colistin Sulphate USP 1,200000 IU Excipients: q.s.	Gastrointestinal, respiratory and urinary-tract infections in calves, goats, poultry, sheep and swine caused by micro-organisms, like Campylobacter, Clostridium, Corynebacterium, E.coli, Erysipelothrix, Haemophilus, Pasteurella, Salmonella, Penicillinase Staphylococcus and Streptococcus spp.	For Oral Administration: Poultry and Swine : 1 kg. per 1000-2000 litre drinking water for 3- 5 days. Calves, Goats and Sheep : Twice daily 5 gm per 100 kg. body weight for 3-5 days. Note : For Pre-ruminant calves, lambs and kids only.	100 gms, 1 kg pouch Pack
HEALTHY CHIC	Lincomycin 11% W/W Animal Feed Premix	Each kg of Healthy Chic contains : Lincomycin base 110 gms	To Improve feed conversion To increase Growth For control of necrotic enteritis	20 gms to 4 gms of Healthy chic/ton of feed	1 kg Pouch 25 kg HDPE Laminated Bag
SURESPIR	Spiramysin Hydrochloride 20%	Spiramycin as a base 20%	Prevention and treatment of Coccidiosis caused by Eimeria sp. in poultry, calves, lambs, kids and rabbits.	Calves : 20g/kg of milk replacer of fedd for 4-5 days Swine : 1 g/ kg of feed for 4-5 days	1 kg pouch , 25 kg HDPE laminated bag
NEOFRED POWDER	Neomycin Sulphate 100 mg	Each gm of Neofred Powder contains: Neomycin Sulphat BP 143 mg Equivalent to Neomycin 100 mg	For The Treatment And Control Of Colibacillosis (bacterial Enteritis) Caused By Escherichia Coli Susceptible to Neomycin Sulphate In Cattle Swine, Sheep And Goats. For The Control of Mortality Associated With Escherichia Coli Organisms Susceptible To Neomycin Sulphate In Growing Turkeys.	Pigs Weighing 25 to 50 Pounds 2 Packets Pigs Weighing 50 to 100 Pounds 3 Packets Pigs Weighing Over 100 Pounds 4 Packets	1 kg pouch 25 kg HDPE laminated bag

ANTICOCCIDIALS / COCCIDIOSTATS

AMPRO -20	Amprolium HCl BP vet 20% w/w Soluble Powder	Each gm of Ampro 20 contains : Amprolium Hcl BP Vet 200 mg .	Ampro 20 is indicated for coccidiosis caused by coccidia susceptible to amprolium (Eimeria spp.), or gastrointestinal infections for which it is therapeutically or prophylactically indicated to administer amprolium, in calves, goats, sheep and poultry.	For oral administration : Calves, Sheep and Goats: Preventive : 1 gram per 50 -100 kg ., body weight through drinking water or milk for 21 days. Curative : 5 gram per 25-50 kg ., body weight through drinking water or milk for 5 days. Poultry : 20 gram per 20-40 litre drinking water for 5-7 days.	1 kg, 25 kg Drum
AMPRO -30	Amprolium HCl BP vet 30% w/w Soluble Powder	Each gm of Ampro 30 contains : Amprolium Hcl BP Vet 300 mg .	Ampro-30 is indicated for coccidiosis caused by coccidia susceptible to amprolium (Eimeria spp.), or gastrointestinal infections for which it is therapeutically or prophylactically indicated to administer amprolium, in calves, goats, sheep and poultry.	Calves, Goats, Sheep: Preventive :1 g per 60 kg body weight through drinking water or milk for 21 days. Curative :1 g per 30 kg body weight for 5 days. Poultry: Preventive :1 kg per 5000 litre drinking water for 1-2 weeks. Curative : 1 kg per 1250-2500 litre drinking water for 5-7 days.	1 kg, 25 kg Drum
TOZU	Toltrazuril 2.5% W/V Oral Solution	Each 100 ml contains: Toltrazuril 2.5 g .	For the control of Coccidiosis caused by : Chicken: E.acervulina, E.brunetti, E. necatrix, E.taenella, Turkey: E. adenoides, E.meleagridis, E.meleagrimitis, Geese : E.anesers, E.truncata, It acts against all intracellular development stages of Coccidia..	28 ml of TOZU per 100 kg body weight per day for 2 consecutive days Equivalent to 7 mg Toltrazuril/ kg body weight per day for 2 consecutive days. Or as advised by the Veterinarian.	100 ml bottle 1 Ltr HDPE Bottle 25 Ltr Drum

NO-COCCI	Maduramycin Ammonium 1% w/w Premix	Each kg of No-Cocci contains: Madramycin activity 10 gms	To control and prevent Coccidiosis in broilers and in growers upto 16 weeks	500 gms per ton of feed	1 kg, 25 kg HDPE laminated bag
DINI 25	Dinitolmide 25 % w/w Animal feed premix	Each 500 gms of (Dini 25) contains: Dinitolmide (Dot) .. 125 gms	For the control of Coccidiosis caused by Elimeria species in layer birds.	For Layers: 0-8 weeks 500 gms / ton of feed. 9-14 weeks 300 gms / ton of feed.	25 kg HDPE Laminated bag.

ANTHELMINTIC

VORMIFRED -OS 2.5	Albendazole Oral Suspension 2.5% w/v	Each ml contains: Albendazole :25 mg	Prophylaxis and treatment of worm infections in Calves,Cattle, Goats and Sheep like: Gastrointestinal worms : Bunostomum, Cooperia, Chabertia Haemonchus, Nematodirus, Oesophagostomum, Ostertagia, Strongyloides and Trichostrongylus spp. Lung worms : Dictyocaulus viviparus and D. filaria Tapeworms : Moniezia spp. Adult Fasciola hepatica.	Goats and sheep: 1 ml. per 5 kg. body weight. Liver-fluke : 1 ml. per 3 kg. body weight Calves and cattle: 1 ml. per 3 kg. body weight.	500 ml & 1 Litre
VORMIFRED -OS 10	Albendazole Oral Suspension 10% w/v	Each ml contains: Albendazole 100 mg	Prophylaxis and treatment of worm infections in Calves,Cattle, Goats and Sheep like: Gastrointestinal worms : Bunostomum, Cooperia, Chabertia, Haemonchus, Nematodirus, Oesophagostomum, Ostertagia, Strongyloides and Trichostrongylus spp. Lung worms : Dictyocaulus viviparus and D. filaria Tapeworms : Moniezia spp. Liver-fluke : Adult Fasciola Hepatica.	For oral administration: Goats and sheep : 1 ml. per 20 kg. body weight. Liver-fluke : 1 ml. per 12 kg. body weight Calves and cattle : 1 ml. per 12 kg. body weight Liver-fluke : 1 ml. per 10 kg. body weight.	500 ml & 1 Litre

PHYTASE

NUTRITASE 2500	Phytase-2500	Each g contains: Phytase 2500 PPU	Improves nutrient digestibility of pH, Amino acids, Protein & Trace Minerals. Reduce accumulation of P in environment. Reduces excretion of nitrogen. Lowers Feed cost.	Layers: 200 gms to 250 gms / ton for replacement of 10 to 12 kgs of Di-calcium phosphate. Broilers: 400 gms to 450 gms / ton for replacement of 10 to 12 kgs of Di-calcium phosphate.	1 kg, 5 kgs , 10 kgs & 25 kg Bag
NUTRITASE 5000	Phytase -5000	Each g contains: Phytase 5000 PPU	Improves nutrient digestibility of pH, Amino acids, Protein & Trace Minerals. Reduce accumulation of P in environment. Reduces excretion of nitrogen. Lowers Feed cost. Lowers Feed cost.	Layers: 200 gms to 250 gms / ton for replacement of 10 to 12 kgs of Di-calcium phosphate. Broilers: 400 gms to 450 gms / ton for replacement of 10 to 12 kgs of Di-calcium phosphate.	1 kg, 5 kgs , 10 kgs & 25 kg Bag

ANTIMYCOPLASMAL

TYLOTAR	Tylosin Tartrate Veterinary Oral Powder	Each 100 g contains: Tylosin Tartrate BP 80 g Equivalent to Tylosin Base	It is useful to control CRD It is useful to improve FCR Increases weight gain Improves Fertility Improves Hatchability.	1 gm in 1.5 ltrs of drinking water daily in 2 divided doses for 2 to 3 days.	1 kg Pouch 25 kg Drum
---------	---	--	---	--	--------------------------

LIVER TONIC

LIVUP (Powder)	Nutrition Supplement	Each Kg contains: Tephrosia Purpurea 5 % Cichorium endivia 12 % Eclipta Alba 11 % Picrorhiza Kurroa 15 % Fumaria Officinalis 10 % Andrographis Paniculata 6 % Phyllanthus Niruri 3 % Tinospora Cordifolia 6 % Terminalia Chebula 14 % Ocimum Basilicum 2 % Emblica Officinalis 16 %	Liver upgradation and protection to improve FCR Builds immunity to Aflatoxins Increases weight gain in broilers and egg production in layers & breeders	500 gms per Ton of feed Or, as directed by the veterinarian	500 g container, 25 kg Pack
----------------	----------------------	--	---	--	-----------------------------

LIVUP (Liquid)	100% Herbal Liver Protector Liquid	Each 100 ml contains: Silybum Mariamum 6 gm Boerhaavia Diffusa 4 gm Tephrosia Purpurea 3 gm Andrographis Paniculata 8 gm Solanum Nigrum 4 gm Terminalia Chebula 3 gm Emblica Officinalis 3 gm Eclipta Alba 2 gm Trachyspermum Ammi 1 gm Zingiber Officinalis 1 gm Carrier q.s.	Improves FCR Resists Aflatoxicosis Overcomes initial resorptive jaundice Liver Protectant and growth promoter Reduces mortality rate Improves liver functions Increase milk production in Mammals and Egg output in Poultry	Horses, Camels & Cows: 40 ml twice a day Calf, Foal, Colt 20-25 ml twice a day Sheep & Goat 10-15 ml twice a day POULTRY: Chicks 5 ml/100 birds per day Growers 10 ml/100 birds per day Broilers 5-10 ml/100 birds per day Layers 15 ml/100 birds per day Or, as directed by the veterinarian.	1 Ltr. & 5 Ltr. Pack
DEXTROVIT	Liquid Feed Supplement	Each 10 ml contains: Choline Chloride : 2000 mg Liver Extract : 125 mg Yeast Extract : 40 mg Vitamin B1 : 2.5 mg Vitamin B12 : 0.75 mcg Niacin : 24 mg DL- Panthenol : 2.5 mg Inositol : 35 mg Base : q.s.	Addition rate (through drinking water) for 100 birds daily : Broilers : 5 - 10 ml/ 100 birds Layers : 20 ml / 100 birds	Large Animals : 30 ml per day Small Animals : 15 ml per day Broiler : 5-10 ml/100 birds daily Layers : 20 ml/100 birds daily Or, As directed by the veterinarian	1 litre, 5 litre HDPE bottle. 10 litre HDPE drum

OTHER PRODUCTS

VIRGIN - 2	Virginiamycin 2%	Each kg contains: Virginiamycin..... 20 gm	In Layers : better Egg laying & better FCR In Breeders : Uniform Eggs In Feed Mills : Energy & Lower density feeds	Broilers : 500 gms/ ton of feed from day- old till marketing Layers Chicks : 500 gms/ ton of feed from day-old to point of lay Adults Layers : 250 gms/ ton of feed from point of to end of lay Breeders : 1 to 2 kg / ton of feed from point of lay to end of lay	1 kg , 5 kgs & 25 Kg Packs
FLAPOL - 8	Flavophospholipol 8%	Flavophospholipol...80 gm	It is useful for improved FCR and growth promotion in chicken It is useful for administration with animal feeding stuff by "Top Dressing"	Broilers :125-250 gms / tonne of feed Layers : 25 to 62.5 gms / tonne of feed	1 kg &25 kg Packs
ENZYPOWER POWDER	Multiple Enzymes Water Soluble Powder	Each Kg contains: Cellulase 2000000 IU Xylanase 200000 IU Amylase 400000 IU Protease 200 IU Phytase 200 IU Carrier q.s.	Optimizes use of both conventional and nonconventional feed ingredients Makes nutrients available in the intestine Improves utilization of Amino Acids Better feed utilization Improves FCR Less mortality and better weight gain Reduce incidences of diseases	Broilers: 300 to 350 g per tonne of feed. Breeders: 250 to 300 g per tonne of feed. Layers: 200 to 250 g per tonne of feed. Or, as directed by the veterinarian.	1 Kg, 25 Kg Packs
ENZYPOWER LIQUID	Multiple Enzymes Liquid	Each 1 Litre contains: Protease NLT 400,000 IU Cellulase NLT 1,00,000,000 IU Xylanase NLT 1,500,000 IU Lipase NLT 6,500 IU Amylase NLT 2,50,000 IU Pectinase NLT 30,000 IU β-Glucosidase 10,000 U α-Galactosidase 10,000 U Arabinase 7,000 U	Increases availability of polysaccharides and polypeptides Makes nutrients available in the intestine Improves utilization of Amino Acids Better feed utilization Improves FCR Less mortality and better weight gain Reduce incidences of diseases Increases phosphorus availability from phytase Increases calcium retention Increases egg shell quality	1 ml/4 ltr. of Drinking Water Or, as directed by the veterinarian.	1 litre, 5 litre HDPE bottle 25 litre HDPE drum
NEOFUR - V	Furazolidone Powder	Furazolidone 20% Powder	In Broilers increases growth rate and improves weight gain by increasing feed conversion. In layers provides better Egg Production better Hatchability, Lower Mortality Effective against the Salmonella species and other infection of Gastrointestinal tract	Routine Inclusion Level : 250gm of FURA-20% in 1000kg of feed and to be fed to Breeders, Chicks, Growers Layers & Broilers, Checks Grower Layers & Broiler Routinely Week- A – Month Plan – 500gm of FURA-20% in 1000kg of feed to be Fed for one week in Each Month of Layer.	1 kg , 5 kg & 25 kg Packs

CYRO -1	Cyromazine 1% w/w Animal Feed Additive	Cyro -1 premix contains : 1% cyromazine	For building an eco-friendly environmental condition for the birds by controlling fly and larvae	500 gms of Larvozine 1% w/w to be added to each toe of feed	1 kg , 5 kg & 25 kg Packs
VITAMINS LIQUID					
OSSO -D	Vitamin D3 2,00,000 IU / gm	Each gram of Osso-D Contains : Vitamin D3.....200000 IU	Protects from deficiency & rickets development Minimises feed cost Optimises metabolism of calcium & Phosphorus Give better FCR & higher body weight Increases quality of Egg shells Develops musculo-skeletal system	Pullet starter 10 to 15 gms Pullet Grower & Developer 8 to 10 gms Layer 10 to 15 gms Broiler 125 to 15 gms	1 kg , 5 kg & 25 kg Packs
FREDS SEL -E (LIQUID)	VITAMIN E WITH SELENIUM ORAL LIQUID	Selenium.....0.5 mg Vitamin E.....100 mg	Fights Muscular dystrophy in broilers and layers Enhance egg production Builds better immune system in birds	2-3 ml in drinking water per 100 birds for all age groups Or, as directed by the veterinarian	1 kg 5 kgs & 25 kg Packs
FRETONE-V	Vitamin A + D3 +E+ C Oral Liquid	Each 5 ml contains: Vitamin A 250,000 IU Vitamin D3 25,000 IU Vitamin E 150 mg Vitamin C 500mg	An anti stress and anti oxidant feed supplement Enhances egg production and hatchability promotes weight gain in boilers	Broiler: 20-50 ml/1000 birds in drinking water. Layers: 30-70 ml/1000 birds in drinking water. Prepare fresh drinking water with Fretone – V daily Or, as directed by veterinarian	1 ltr , 5 litre & 10 litre Packs
FREDS	Vitamin B Complex + DL-Methionine Liquid	Vitamin B2 2 mg Vitamin B6 0.62 mg Vitamin B12 6.25 mcg D-Panthenol 1.25 mg Niacinamide 37.5 mg DL-Methionine 5 mg L-Lysine 5 mg Choline Chloride 5 mg	A Vitamin Supplementation to birds for better performance.	Broilers :2-5ml/100 birds in drinking water Layers :3-6ml/100 birds in drinking water Or, as directed by the veterinarian.	1 ltr 5, ltr & 10 Ltrs Packs
FREDCAL -D	Liquid Animal Feed Supplement of Calcium, Phosphorus, Vitamin B12 & Vitamin D3	Each 20 ml of Fredcal-D contains: Calcium.....325.6 mg Phosphorus.....167.7 mg Vitamin B12.....20mcg Vitamin D3.....1600 IU Flavoured Base q.s.	Optimises Egg quality Improves quality of egg shell Prevents rickets in birds For prevention of prolapse/cannibalism.	Cattle & Horse :100ml daily Calves :40 ml daily Dogs : 20 ml daily Poultry : 20 ml daily for 100 birds Broiler & Growers:50 ml daily fir 100 birds Layers : 100 ml daily for 100 birds Or As directed by the Veterinarian	5 ltrs , 5 ltr & 25 ltrs Packs
FREDCAL -D PLUS	Liquid Animal Feed Supplement of Calcium , Phosphorus, Vitamin D3, B12 along with Herbal Extracts	Each 100 ml contains: Calcium Phosphate 7.5 gm (Elemented Calcium) 2.2 gm (Elemented Phosphorus) 1.7 gm Vit B12- 100 mcg Vit D3- 10,000 I.U. Aqueous extract derived from Jivanti 4 gm Shatavari 8 gm Jatamansi 1 gm Kalongi 0.6 gm	Calcium Phosphorus : Provide high quantities of calcium and phosphorus in the ratio of 12:1 which matches the ratio required by lactating cow and buffalo Vit D3 : ensures maximum absorption of calcium and phosphorus from the feed Vit D12 : maintains appetite and is also necessary for metabolism of Propionic acid in liver Jivanti : A potent galactagogue increases milk production Shatavari : A potent galactagogue with immunomodulant action, giving high immune status and improves milk production kalongi : increases number of milk secreting cells in udder, thus increases milk producing capacity of udder Jatamansi : Reduces restless-ness thus assures better letting down of milk by reducing anxiety during milking and helps in complets milking.	Oral - Direct or in Feed Cattle/ Buffalo : 40 ml twice daily for 12-13 days before and 12-13 days every month Dog : 5 ml twice daily	1 Ltr , 5 Ltr Packs
VITAFRED	Vitamin A solution	Each ml of Vita powder contains: Vitamin A.....100000 IU Excipients q.s.	Increases egg production and hatchability Builds resistance in Enteritis, Coccidiosis, Infections bronchitis. Prevents Blood spots in the egg. Supplement therapy during CRD Coryza and helminthiasis.	To be given at least 1 week in a month. Broiler : 2 ml / 100 birds Layers : Chicks : 2 ml / 100 birds Adults : 6 ml / 100 birds	1 ltr , 5 ltr & 25 ltr Packs
MULTIMINERALS & MULTIVITAMINS POWDER & PREMIX					

FREDS- BE	Vitamin B Complex Feed Premix four strength	Each kg of Freds-Be contains: Vitamin B1.....16 gms Vitamin B6.....32 gms Vitamin B12.....0.16 gms Vitamin E..... 160 gms Calcium Pantothenate 160 mg Niacin.....340 mg Folic Acid.....16 mg L-Lysine.....40 mg DL-Methionine.....40 mg Calcium......52 mg Carrier.....q,s	Buils resistance to fight against diseases Important for productions and utilization of energy Provides vitamins required for growth and production.	50 gms of Freds BE per ton of feed	1, kg, 5 kg , 10 Packs
C-MAGIC	Water Misscible Vitamin C 100% Feed Supplement Powder.	Each gram of C-Magic contains: Vitamin C 1000 mg	C-Magic is an essential vitamin for calves,cattle, goats, poultry, sheep and swine. C-Magic is used for: Prevention or treatment of vitamin C deficiencies in farm animals Prevention or treatment of stress (caused by vaccination, diseases transport, high humidity, high temperatures or extreme temperature changes	For oral administration Poultry, swine : 100 gram per 2000 litre drinking water for 3 - 5 days. Calves, goats and sheep : gram per 200 kg. body weight for 3 - 5 days. Cattle : 1 gram per 400 kg. body weight for 3 - 5 days.	Sachet of 100 gram and jar of 500 & 1 Litre.
CELECTO	Electrolyte Powder with Vitamin C	Each 100 g contains : Sodium Chloride 1 gm Potassium Chloride 3 gm Sodium Carbonate 3 gm Sodium Acid Phosphate 0.008 gm Calcium Lactate 1.1 gm Magnesium Sulphate 1 gm Calcium Gluconate 1.1 gm Vitamin C 1 gm Carrier q.s.	To avoid dehydration & heat strokes in summer.Useful in Diarrhoea and To maintain perfect osmotic pressure and precise acid base equilibrium	General Usage & Adminstration Through feed : 1 kg per ton feed Through Drinking water Chicks : 100 gm per 1000 chicks Growers : 75 gm per 500 birds Layers & Broilers : 1 gm per 2 ltr of water Breeders : 1 gm per 1 ltr of water	1 kg , 5 kg
OSSITONE GRANULES	Biologically derived Calcium Phosphorus Feed Supplement with Trace minerals & fortified with Vitamin D3	Each 5 g contains : Biologically derived Micro Crystalline Hydroxyapatite Compound (MCHC) equivalent to Calcium : 400 mg Phosphorus : 190 mg with trace minerals like Copper, Cobalt, Iron,Magnesium and Collagenous proteins Fortified with Vitamin D3 : 1500 IU	OSSITONE Granules is a unique blend of Microcrystalline Hydroxyapatite Complex (HPMC) a biologically derived calcium, phosphorus with Vitamin D3 The MCHC improves Skeletal and muscular strength in animals Prevents osteoporosis and rickets. Whereas, Vitamin D3 regulates calcium and phosphate metabolism, guaranteeing calcium absorption by the intestinal mucosa	Normal mixing ratio : 250 g – 500 g per ton of feed Cattle & Horses: 30 g - 50 g daily Sheep, Goats & Pigs: 10 g Dogs & Cats: 5 g - 10 g To be given daily for 10 days continuously in a month or as directed by the Veterinarian.	250 g & 500 g Jar packs laminated bags
LAYERS PREMIX PLUS		Each kg of Layers Premix Plus contains Vitamin A 90,00,000 IU Vitamin D3 21,35,000 IU Vitamin E 20,000 mg Vitamin B1 1340 mg Vitamin B2 5340 mg Vitamin B6 2500 mg Vitamin B12 25 mcg Folic Acid 1000 mg Manganese 117340 mg Zinc 78000 mg Iron 63340 mg Copper 10670 mg Potassium 10000 mg Calcium 20000 mg Niacin 30000 mg Partially hydrolyted proteins 300 gm* Casein having a balanced Amino acids profile as follows: Lysine 7.2% Gutaric Acid 17.75% Cystin 0.40% Valine 6.55% Tyrosine 5.44% Aspartic Acid 5.95% Histidine 2.72% Proline 11.63%	Highly essential for the improved animal's performance and health Provides enhanced digestibility of the feed materials, Improves the quality of feed	Inclusion Rate : 200 - 500 g per ton of feed	1 kg , 5 kg & 25 Kg HDPE Laminated Bag

BROILER/BREEDER PREMIX PLUS		<p>Serine 5.78% Tryptophane 1.40% Arginine 3.57% Threonine 3.83% Glycine 1.79% Alanine 2.81% Methionine 2.98% Isoleucine 6.38% Leucine 8.50% Phenylalanine 5.6% Appropriate overages added for losses on storage.</p>	<p>Delayed growth, Reduces hatchability of eggs Loss of appetite For feathering in birds Swelling of eyelids Fatty hypertrophy of liver</p>	Inclusion Rate: 200 gm per tonne of feed	<p>1 kg, 5 Kg 25 kg HDPE Laminated Bag</p>
FREDS SEL -E	Vitamin E and Selenium Premix	<p>Each kg of Freds sel-e contains: Vitamin E.....20%w/w Selenium.....400 PPM</p>	<p>A supplement for Vitamin E deficiency To increase productivity To combat muscular dystrophy</p>	500 gm / ton of feed	<p>1 kg x 10 jars shippers 25 kg HDPE laminated bags.</p>
FREDMIX	Vitamin AB2D3K Feed Premix Triple Strength	<p>Each gram of Fredmix Contains: Vitamin A.....2,47,500 IU Vitamin B2.....150 mg Vitamin D3.....3600 IU Vitamin K.....30 mg</p>	<p>Improves feed conversion Improves egg production in layers Improves weight gain in broilers Improves growth of chicks and reduces mortality Prevents curled toe paralysis Ensures proper coagulation of blood</p>	35 to 40 gms per ton feed	<p>1 Kg , 5 Kg & 25 kg HDPE laminated BAGS</p>
OSSIPOWER	A unique blend of Glucosamine Sulphate, Methyl Sulphonyl Methane, Chondroitin Sulphate, Boswellic Acid with Vitamins (Ascorbic Acid), Tocopheryl Acetate, Minerals (Manganese Chloride & Zinc Chloride) & Natural source of Calcium	<p>Each 5 gm powder contains: Glucosamine Sulphate USP 400 mg Methyl Sulphonyl Methane 100 mg Chondroitin Sulphate 20 mg Ascorbic Acid 100 mg Tocopheryl Acetate 50 % with Vitamin Acetate 10 mg Manganese Chloride 0.50 mg Zinc Chloride 0.50 mg Boswellic Acid 45 mg Microcrystalline Hydroxyapatite q.s. Complex (MCHC)</p>	<p>For reducing inflammation & pain associated with acute inflammation of body parts of a animal, particularly joints due to injury or due to arthritic conditions like osteoarthritis and rheumatoid arthritis or other disease.</p>	<p>5 g per day with or after meals for small animals & 5 g twice a day with or after meals for large animals. Allow 4 – 6 weeks for best results If lameness worsens or persists, discontinue use and contact your veterinarian.</p>	<p>250 g & 500 g Jar packs laminated bags</p>
OSSITONE +	Combination of Biologically derived Calcium Phosphorous feed supplement with potent herbal extracts	<p>Each 50 g contains: Microcrystalline Hydroxyapatite Complex (MCHC) providing Calcium 9 gm Phosphorous 4 gm With trace minerals like copper, cobalt, iron, magnesium Fortified with 12.5 gm mixture of herbal extracts containing: Shatavari, Jivanti, Methi, Galo, Ashwagandha, Jatamashi, Varahikand, Vidarikand, Piper, Piparimul</p>	<p>OSSITONE+ is a unique blend of Microcrystalline Hydroxyapatite Complex (HPMC) a biologically derived calcium, phosphorus with potent galactagogue herbal extracts. The MCHC improves 1. Skeletal and muscular strength in animals 2. Prevents osteoporosis and rickets. Whereas, the herbal extracts like Shatavari, Jivanti, Methi, Galo, Ashwagandha, Jatamashi, Varahikand, Vidarikand, Piper,</p>	Add daily OSSITONE+ (50 g) in feed of each animal	<p>50 g Sachet pack 100 g & 500 g Jar pack</p>

IMMU – 365	Immuniser & Stress Reliever activation, building of immunity, and stress relief.	Each Liter contains: Aswagandha (Leaves) 10,000 mg Bhrami (Roots) 20,000 mg Chitrak 15,000 mg Camellia Sinensis 10,000 mg Phyllanthus Emblica 20,000 mg	Piparimul are combined to strengthen this formula. This herbal mixture 3. Improves milk yield and helps in irregular or suppressed lactation in cattle and buffaloes. 4. Ensures lengthy lactation. 3. Prevents post-partum haemorrhage. 4. Treats psychological Agalactia due to death of calf or first lactation or change in milker. New born chicks Vaccinations Debeaking, Deworming Climatic disturbances, Transportation Poor growths, Uneven sizes Disease conditions, Lower / Higher egg productions etc.	1 Liter Immu-365 per 500 kgs. of feed or 1 Liter Immu-365 per 15,000- 20,000 Kgs. body weight or	1 Ltr , 5 Ltrs & 10 Ltrs. Packs
PROCHIC	Multi Strain Probiotic for Poultry & Animals	Each gram of PROCHIC contains: Bifidobacterium bifidum 2.00*10 ⁸ cfu/g Lactobacillus acidophilus 2.06*10 ⁸ cfu/g Lactobacillus bulgaricus 2.06*10 ⁸ cfu/g Lactobacillus casei 2.06*10 ⁸ cfu/g Lactobacillus plantarum 1.26*10 ⁸ cfu/g Streptococcus thermophilus 4.10*10 ⁸ cfu/g Yeast Torulopsis spp 5.32*10 ⁷ cfu/g Aspergillus oryzae 5.32*10 ⁷ cfu/g Total Viable Count 2*10 ⁹ cfu/g	Reduces digestive disorders. Improves feed conversion ratio (FCR). Improves egg shell quality. Improves food safety. Helps in longer laying life. No withdrawal period is required.	DOSAGE IN WATER: For Poultry: Chicks: 1 Gram per liter of water for a period of 5-7 days. For Poultry: After antibiotic therapy: 1 Gram in 4 liters of water for 5-7 days. For Poultry: During stress conditions: 1 Gm in 4 liters of water for 3 days. DOSAGE IN FEED: For Poultry: Broiler: Pre-starter: 150 Grams per ton of feed. For Poultry: Starter: 100 Grams per ton feed. For Poultry: Finisher: 50 Grams per ton of feed. Use for a For Poultry: Breeder & Layers: Use 50 to 100 Grams per ton Can also be given to big animals. 2-3 Gms for 10Kg body weight or as directed by veterinarian.	1 kg , 5 kg & 10 kgs Pack

DISINFECTANTS

SANIFACT	Water Sanitizer for all types of Live Stock	A blend of Peracetic Acid Hydrogen Peroxide. Acetic Acid Sulphuric Acid and Surfactant in a stabilized aqueous solution	For Sanitizing Water: Use SANIFACT 5- 10 ml per 100 liter of water. In acute cases: Use 15-20 ml. of SANIFACT in 100 Ltr. of water. In severe cases: Use 40- 50 ml of SANIFACT per 100 Ltr. of water can be continued for 2 -3 days. For Farm Sanitization with SANIFACT: Empty Sheds/Barns, Sanitization of floor and wooden or concrete surface For Cleaning and Sanitization of Equipments, Footbath / Wheel dip For Pipeline Cleaning (Pipes must be of non-corrosive plastic material) Hatchery Disinfection with SANIFACT: On Floor, Humidity Chamber, Eggs Before Hatching Incubator Tank , Hatchery/Setter (Floor and Trays)	Sanifact is useful for control of recurrent bacterial and viral infections present in all water sources.	5 Litre, 15 Litre 25 litre drum container
FREQUAT	Water Sanitizer And Utility Equipment Cleaning Agent	A unique Synchronised Blend of Quaternary Ammonium Compounds Pine oil. Citronella oil and Surfactants. IPA in aqueous base.	Kills gram negative and gram positive bacteria, molds and viruses EPA registered to sanitize vacant animal by thermal fogging Leaves a residual bacteriostatic film on treated surfaces Economical use dilutions (½ ounce per gallon or 4 ml/ liter to disinfect) Safe to use on most surfaces not harmed by water Useful for manual cleaning and for foam cleaning Fogging hatchers, setters and hatchery rooms	Use 15 ml per 100 litres of water for prevention In other use 20 ml per litre for spray or mopping	1 litre and 5 litre bottle, & 25 litre drum container

LIVE STOCK / BOLUS TABLETS

FENDA-250	Fenbendazole Bolus 250 mg	Each g contains: Fenbendazole BP 250 mg	Cattle, Buffaloes, Sheep and Goats All important adult nematodes, including inhabited Larvae, Lung worm and tape worms Horses : Strongyles, Ascaridis, Oxyuris and Strongyloides Pig : Hyostrongylus rubidus, Oesophagostomum spp., Ascaris suum Trichuris suis, Metastrongylus spp., Stephanurus dentatus (kidney worm) Dogs : Ancylostoma (Hook Worms) Ascaridis, Trichuris spp. (Whip Worm) and Taenia spp	Cattle & Buffalo : 5 mg / kg body weight Sheep & Pigs : 5 mg / kg body weight Horses : 5-10 mg /kg body weight Dogs : 50mg/kg for 5 days	Strip of 4 Boluses 20 Bolus Box 5x4 Bol Alu.Strip
-----------	---------------------------	--	---	---	--

PIQAN	Praziquantel Tablet 50 mg	Each tablet contains : Praziquantel....50 mg	Piqan is active against all adult and development stages of Tapeworms such as Echinococcus granulosus , Echinococcus multilocularis, Taenia hydatigena Taenia multiceps Dipylidium, caninum Mesocostoides sp	It is given as a single dose of 5 mg/kg body weight orally in case of dogs and cats after every six weeks starting from the age of six weeks. The dose schedule is as follows : B.W. - 5 kg : 1/2 Tab B.W. - 10 kg : 1 Tab B.W. - 15 kg : 1 + 1/2 Tab B.W. - 20 kg : 2 Tab Over 20 kg increase dose rate by 1/2 for every 5 kg increase in weight	Blister of 4 tablets
PEFXA	Pefloxacin Tablet 400 mg	Each tablet contains : Pefloxacin.....400 mg	For the treatment of gastrointestinal, respiratory, urinary and genital tract infections. Broad Spectrum Bactericide covering Mycoplasma spp. Also ideal for pre-ruminant calves and monogastric animals Pefxa belongs to Fluroquinolone group of antimicrobials. These compounds are regarded as almost the ideal ant- microbial agents because of their broad spectrum of activity, clinically advantageous pharmacokinetic properties & low toxicity.	1 tablet per 80 kg body wt for 3-4 days	Strip of 4 Tablets
RAF XI	Rafoxidine bolus 2 gm	Each bolus Contains : Rafoxidine 2 gm	Effective against immature and flukes as well as blood sucking nematodes, tissues invading fly maggots and nasal bots (Oestrus ovis). Safety Margin is wide and can be used at therapeutic dosage in cattle and sheep of all ages under commonly encountered conditions without hazards	Basic Dose 7.5 mg/kg body wt. Cattel/ Buffalo : 1 Bolus . In case of maggot infested wounds only repeat after 3 days	Strips of 2 bolus
STANFRE	Clostantel bolus 1000 mg	Each Clostan bolus contains : Clostantel.....1000 mg	Clostantel, a long acting ectendoparasiticide (effective against both ecto & endoparasites) is highly effective against the following endoparasites & ectoparasites in cattle, buffalo, sheep, goat, dog, horse and other domestic animals. Endoparasites : Fasciola hepatica, fasciola gigantica, Fascioloides magna, Haemonchus contortus, Haemonchus similis, Bunostomum phlebotomum Oesophagostomum radiatum, Strongyloid spp, Ascarid spp, Ancylostma caninum, Ectoparasites : Oestrus ovis, Hyalomma anatolicum, Boophilus microplus, Dermatobis hominis, Ligonathus ovillus, Callitroga hominivorax Melophagus ovis, Gastrophilus spp, Lucilia cuprina Schistosoma nasalis	For Fasciola hepatica : Fasciola gigantica, Fasciola gigantica, Fasciola magna : Clostan Bolus : 1 Bolus / 66 kg B.Wt For Haemonchus spp. Bunostomum phlebotomum Oesophagostomum radiatum Strongyloids spp. Ascarid spp, Ancylostoma caninum Stanfre : 1 Bolus / 100 kg B.Wt.	Strip of 4 Boluses
HALIFRED	Halquinol bolus 1.5 g	Each bolus contains : Halquinol 1.5g	For treatment of diarrhoea	1/4 - 1/2 bolus twice daily orally for 5-7 days	Strip of 4 Boluses
VORMIFRED -V	Albendazole bolus 1.5 g	Each bolus contains: Albendazole U.S.P....1.5gm	A true broad spectrum anthelmintic with high efficacy against immature and mature gastrointestinal roundworm Lungworm, tapeworm, liver fluke and amphistomes in ruminants	For Round Worms Cattle / Horse : Bolus / 300 kg B.W. For Fukes : Cattle : 2 Bolus per 300 kg. B.W	20 Bolus Box 5x4 Bol Alu.Strip
MEBENFRED -V	Mebendazole bolus 1 gm	Each bolus contains : Mebendazole.....1 gm	Effectively kills ova, larva, immature and mature stages of Roundworms & Tapeworms in gastrointestinal & respiratory tract without causing any stress Equally effective in all species of animals Cattle / Sheep : Haemonchus (Barberspole), Ostertagia, Trichostrongylus Nematodirus, Bunostomum (Hookworms), Oesophagostomum Dictyocaulus, (Lungworms), Chabelia, Cooperia, Cestodes (Tapeworms). Horse : Large and small strongyles Parascaris (Roundworms) Oxyuris equi (itch and Pin Worm) Tapeworms Probstmayria, Vivipara	Cattle 4-5 bolus Sheep/ Goat 1-2 bolus Horse 2-4 bolus	Strips of 4 bolus
DIMIFRED	Diminazene Diacetate Granules	23.6 g of granules containing: 10.5 g of the active Ingredient	Diminazene is highly active against Trypanosoma spp, Babesia spp, and Theileria annulata	Each sachet contains: 10.5 g diminazene diacetate sufficient for 3000 kg of body weight. To be dissolved in 150ml of water for the preparation of an injectable solution	23.6 gm Sachet 2.36 gm Sachet
PRAZFEB	Praziquantel tablet 50mg & Febendazole tablet 500 mg	Each tablet contains : Praziquantel U.S.P.....50mg Febendazole B.P vet 500 mg	For regular deworming of tape worms, Praziquantel kills mature & immature stages of common species of tapeworms like Echinococcus granulosus, E-multilocularis, Taeniainmulticeps, T.Hydatigena, T.Pisiformis, T.taeniformis Dipylidium caninum.	One Prazfeb tablet for 10 kg body weight of the dog as a single dose Prazfeb tablet @ 5 mg/kg body weight and Febendazole @50 mg / kg body weight The dose is to be repeated every month.	Strip of 2 tablets in a catch cover

FREMET-V	Metronidazole tablet 200 mg & 400 mg	Each tablet contains: Metronidazole 200 mg & 400 mg	For regular deworming of roundworms Febendazole is very effective on ascardis (Toxocara canis, Taxascaris Leonina), hookworms (Ancylostoma caninum), hookworms (Ancylostoma caninum), Uncinaria, Stenocephala and Whipworms (Trichuris vulpis). Febendazole works on both mature and immature stages of namatodes . Prazfeb tablets is to be given to pregnant bitches & puppies.		
COTRIFRED -V	Trimethoprim tablet 400 mg & Sulphamethoxazole 2 gm	Each bolus contains: Trimethoprim.....400 mg Sulphamethoxazole.....2 g	In enteritis, diarrhoea, dysentery.		
OXY-TRACY	Oxytetracycline tablet 500 mg	Each tablet contains : Oxytetracycline HCl...500 mg	Diarrhoea caused by oxytetracycline sensitive bacteria Prolonged oral administraton may disturb the intestinal- microflora resulting in avitaminosis and infection with candida.		
FREMET-F	Metronidazole tablet 1 g & Furazolidone 200 mg	Each bolus contains : Metronidazole.....1 g Furazolidone : 200g	For treatment of gastrointestinal infection due to G+ve, G-ve aerobic and anaerobic bacteria and protozoa		
TERAFRED	Teramisole Powder	Each kg bolus contains: Tetramizole HCl B.P. Vet 30% W/W	Broad Spectrum anthelminitic for Cattle, Buffalo, Sheep, Goat and Poultry		
PRIME -CAL	Combination of Natural Calcium, Phosphorus & Vitamin D3 Tablets with 25% Proteins, 12% Collagens and Trace minerals	Each tablet contains : Microcrystalline Hydroxyapatite Complex (MCHC) providing Calcium : 500 mg Phosphorous : 220 mg Vitamin D ₃ : 400 IU Proteins : 25% (Approx) Collagens : 12% (Approx) Trace Minerals – Copper, Zince, Magnesium Iron & Amino acids	Prime – cal tablets are recommended as a daily supplement for dogs and cats where appropriate to provide a dietary source of calcium, phosphorous, and Vitamin D.		
				20 mg per kg body weight daily orally for 5-7 days.	Strip of 10 tablets
				Dog and Cat : 30 mg per kg body weight daily orally for 3- 5 days	
				Dog : 1 tab per 10 kg body weight given orally	Pack of 10 tablets
				1/4 to 1/2 bolus twice daily orally for five days	Pack of 10 tablets
				Large Animals: Dissolve 10 gms in 300 ml water and administer in following dose Upto 15 kg Body weight -15 ml 15 to 25 kg Body weight - 30 ml Over 35 kg Body weight - 60 ml Poultry : Dissolve 10 gms powder in 1 ltr. Drinking water for 100 birds	100 gms containers & 10 gms pouch
				Dogs and Cats : 1 tablet per 20 lbof body weight daily.	200 tablets in a Jar
				Pups : 1 tablet BID Small breed dogs : 1 tablet BID Large breed dogs : 2 tablets BID Administer by hand just prior to feeding, or crumble and mix with food.	